

Annual Congress on

Mycology and Fungal Infections

November 16-17, 2017 Atlanta, Georgia, USA

Draft genome sequence of 11 clinical and environmental Colombian isolates of genus *Aspergillus*

Gomez OM^{1,2}, Freyle C G^{2,3}, Tamayo D P^{2,3}, Rua A L^{2,3}, McEwen J G^{2,4} and Hernandez O^{2,3}¹Universidad de Antioquia, Colombia²Corporación Para Investigaciones Biológicas, Colombia³Universidad de Antioquia, Colombia⁴Universidad de Antioquia, Colombia

Background: The genus *Aspergillus* is a group of opportunistic fungi that cause infections, with high morbidity in immunosuppressed patients. Approximately 350 species have been described in this genus, classified in 4 subgenus and 20 sections. In Colombia, *Aspergillus fumigatus* is the most frequent species in these infections. However, in the last years an increase in the incidence of other species has been observed. This added to an increase in the quantity and diversity of *Aspergillus* isolates in hospital environments.

Aims: Characterize the phenotype and genotype of clinical and environmental Colombian isolates of *Aspergillus* spp.

Methods: We collected 11 Colombian isolates of *Aspergillus*: 4 from different clinical samples, six from hospital environments and 1 from extra-hospital environments. The isolates were identified according to their macro and microscopic characteristics. Genomic DNA for sequencing was prepared from mycelium culture using phenol/chloroform extraction. Library preparation and 150-bp paired-end sequencing was performed using the Illumina HiSeq 2500 platform. The reads were de novo assembled using SPAdes 3.10 pipeline. The draft genome assembly quality was analyzed by QUAST. Augustus v3.0.1 and GlimmerHMM were used for gene prediction. Inference by sequence homology was performed with OrthoFinder v2.0.9. IQtree v1.4.4 software was used for phylogenetic reconstruction by ML.

Results: In this study, four of the eleven isolates classified phenotypically were re-classified correctly after genome analysis. We obtained whole genome sequence of 4 clinical and 7 environmental isolates, including 3 novel genome sequences of the genus *Aspergillus* not reported in the databases: *A. tritici* (Clinical), *A. tamari* (hospital environment) and *A. amoenus* (extra-hospital environment).

Conclusion: The results of this study provide useful data for the genomic comparison between clinical and environmental isolates of *Aspergillus*. Valuable information that can be used to understand the evolutionary relationships, determine the diversity and level of production of enzymes and secondary metabolites, predict the pathogenicity and response of this opportunistic fungus to antifungal agents.

Biography

Gomez OM is currently working as Faculty at Universidad de Antioquia, Colombia.

oscargomez24@hotmail.com

Notes: