

# Anthocyanins: Ethnomedicine to Clinical Trials

Komal Pandey<sup>1</sup>, Ashutosh Goswami<sup>1</sup>, Khemraj Bairwa<sup>1</sup>, Abhijeet S. Kate<sup>1</sup>, Debanjan Chatterjee<sup>2\*</sup>

**ABSTRACT:** Anthocyanins are the secondary metabolites chiefly present in the plant cell sap and vacuole. These are mainly responsible for attractive colors in leaves, flowers, and fruits. In the human being, anthocyanins are well known for their beneficial therapeutic effects such as prevention of cancer, oxidative stress, diabetes, obesity, etc. Owing to their huge therapeutic potentials, anthocyanins are highly consumed as nutraceuticals form, herbal medicine, and functional foods. Chemically anthocyanins are flavylium cation linked with phenolic hydroxyl, methoxyl, and sugars groups. Therefore they are also known as polyphenolics with highly aqueous solubility. The sugar-free part of anthocyanins is known as anthocyanidin. Based on the position and number of hydroxyl and methyl groups, anthocyanins were classified in several types. Till now, more than five hundred anthocyanins have been explored which are classified in nine groups and fifty-nine subgroups. The pelargonidin, cyanidin, delphinidin, petunidin, peonidin, and malvidin are the commonly occurring anthocyanins in nature. Anthocyanins have a hydrolysable group and reactive scaffold

which can easily act with the target protein and give plausible interaction. It also has the reactive oxygen species scavenging activity and hence acts as a potent antioxidant effect. Furthermore, anthocyanins are much explored as anti-obesity, antioxidant, anti-diabetic, and anti-cancer agents. Some anthocyanins are in phase 1 and phase 2 clinical trials for diabetes and obesity treatment. There are several world patents, united states patent, Chinese patents, and Korean patents on anthocyanins; particularly in the anti-diabetic area. Overall, the anthocyanins have been highly explored and are the high interest of molecules, but, there is no comprehensive review available for the scientific community which provides detailed phytochemistry, spectroscopic analysis, biosynthesis, biological activities, clinical status and patent-related information of anthocyanin. Therefore, in the present review, the authors tried to fill this gap by summarizing all the scientific information and researches on anthocyanins.

**Keywords:** Anthocyanins, Polyphenones, Anticancer activity, Antioxidant activity, clinical status.

## INTRODUCTION

Healing, through nutritional therapy, is a pretty vintage concept. Since the inception of human civilization is being carried forward and continuously going on, this can give us peace not only our physique but also the soul and spirit (1). In the previous century, natural therapies include too functional foods along with medicinal food (2). The pharmacological effects of plants are mainly associated with secondary metabolites (3).

Anthocyanins are the complex polyphenolic and water-soluble plant pigments, and responsible for attractive pigments mainly in flowers, leaves, and sometime in stem and fruits. Anthocyanin comes from the Greek word “anthos” equals to “Flower” and kyaneos equals to “dark blue” in 1835 (4). Blueberry, raspberry, black rice, and black soybean having red blue-black colors are the few of the richest source of anthocyanin. The anthocyanins are generally synthesized in higher plants by phenylpropanoid pathway. European Union (5).

### Historical of anthocyanin:

In 1664 Boyle in his research color change has been noticed in “syrup of violet” intensified from purple to green which due to “acid liquor”; and the published first book on anthocyanins was “The anthocyanin pigments of plants” by Muriel Wheldale in 1916. Simultaneously, few more scientists were working on anthocyanins including Willstatter and Everest (1913), Willstatter and Nolan (1915), and Shibata. (1919) and concluded the depending upon the pH changes of anthocyanin colors of *Centaurea cyanus* flowers. They also mentioned the first time that the development of *C. cyanus* petal color is due to the conjugation of anthocyanins with metal ions (6). The molecular structure of anthocyanins was discovered by Richard Willstatter and co-workers from 1912 to 1916 (7). After this, Pauling 1939 proved the relationship between colour and anthocyanin. He suggested that colors in anthocyanins are due to the resonating structure of the flavylium cation. The colours of anthocyanins sometimes depend on the pH and pKa of the fluids cell sap and also responsible for particular pharmacological effect in human being (8). For instance, French people have a less coronary arterial disease as they used to take red wine enriched with anthocyanin (9).

Anthocyanins are well recognized in the traditional system of medicine including traditional Chinese medicine and Ayurveda. In France, anthocyanin formulations are widely used in motion sickness. In Germany, these are used in peristalsis and for reducing the gastric distress. Additionally, the recent biological studies also suggest the importance of anthocyanin as antiemetic, positive inotropic, and as a potent morning sickness reducer in pregnant women. Chemically, the anthocyanins are flavylium cation linked with phenolic hydroxyl, methoxyl, and sugars groups. The aglycone of anthocyanins is known as anthocyanidin and glucose, arabinose, galactose, and rutinose are the most commonly found glycone in anthocyanins. Based on the number and position of hydroxyl and methyl groups, anthocyanins were mainly sub-classified into seven categories viz Aurantidin, Malvidin, Pelargonidin, Europinidin, Cyanidin, Petunidin, and Rosinidin.

This variety of anthocyanins has several biological effects over various organs or tissues including pancreas, liver, white adipose tissue, skeletal muscle. They may exert their various physiological effects via decreasing the glucose level, reducing the apoptosis, increasing the  $\beta$  cell viability, increasing the glucose uptake by skeletal muscle, and by reducing the insulin resistance. The safety and efficacy of anthocyanins were also evaluated in clinical trials by National Clinical Trials authority especially for anti-obesity, antioxidant and anti-cancer activities with NCL number NCT01245270, NCT01005420, and NCT 01883401 respectively. Additionally, plenty of patents including world patent or country-specific patents have been filed or granted on anthocyanin. Overall, the anthocyanins have been highly explored as potential therapeutic agents especially for diabetes treatment, inflammation, cancer, and an antioxidant. However, there is still a very less comprehensive review available on anthocyanins which describe the detailed phytochemistry, spectroscopic analysis, biosynthesis, therapeutic activities, clinical status, and patent. After conceiving these facts, the authors tried to fill this gap in the present review.

### Chemistry of anthocyanin

Anthocyanidins are the fundamental structures that are aglycones with anthocyanin and it consists of aromatic ring attached heteroaromatic ones, that is with oxygen bonded with C-

<sup>1</sup>Department of Natural Products, National Institute of Pharmaceutical Education and Research- Ahmedabad, Palaj, Gandhinagar-382355, Gujarat, India.

<sup>2</sup>Department of Natural Products, National Institute of Pharmaceutical Education and Research- S.A.S Nagar, 160062, Punjab, India.

Correspondence: Debanjan Chatterjee, Ph.D. Research Scholar, Department of Natural Products, National Institute of Pharmaceutical Education and Research- S.A.S Nagar, 160062, Punjab, India. Tel: +91- 7980568998, Email: debu06781@gmail.com




This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact [reprints@pulsus.com](mailto:reprints@pulsus.com)

C in the 3rd aromatic ring (10). Pauling 1939 shows the color and proved the resonating effect and structure all around literature reported more than five hundred anthocyanins and among them, only 58 subclasses and then that have more or less 23 anthocyanidins with the 6 different have vascular plants (11). These most commons are Pg, Cy, Pn, Dp, Pt, Mv. There are nonmethylated anthocyanidins that is Cy, Dp, Pg maximum in nature. (12) Among the above 82% are pigmented fruits and some of vegetable. Then 69% of fruits almost 49% in flower distribution more than 61%, in Cy 51% Dp 12% Pn 12% Pg 12% Pt % Mv %. The widely spread things 3 monoxides, 3 biocides 3, 5- 3, 7 -glucoside. They are most present in Cy -3- glucoside. Enormous groups present for the natures beauty. the chemotaxonomic and the adulteration variety also found due to their componential analysis and related compound percentage (13) (14).


2 Phenyl benzopyrilium and flavylium salt main derivatives of this hydrolyzable, they are poly hydroxyl poly methoxy of the anthocyanin moiety. The hydrolyzable groups are methylating in nature and they are attached with sugar molecules. Hundreds of molecules are the same category and they have the carbon skeleton (15). Characterization is done based on the spectrophotometric and the spectrometric tools. Natural resources have a large variety over three chiral centres and their varieties this is also has the same. The hydroxyl groups present and the related derivatives there hemiketal and hemiacetal groups also shown a good number of activity in vivo and in vitro (16). If we found the derivatives they are the hydroxyl methoxyl methyl, methoxy ether. Biosynthetic pathways are the same for the ethoxy mether group anthocyanins. The flavonoid class has the main identity of the 3 ring skeleton and so on anthocyanin likewise (17). On a general time, sugar is attached with C-3 and C-5, exceptional case C-7, then glucose arabinose and galactose are also attached to this. Most frequently acetylation gives the sinapic acid, ferulic acid, acidic acid, and para coumaric acid. Acetic acid, malonic, oxalic and malonic acid. Anthocyanins have a natural shielding of aqueous attack. Bathochromic, hypsochromic shifts are also there in UV spectra. According to the variability structure the anthocyanin has fragmentation variety and there isomerization pattern on each and every spectra (18).

#### Ground state chemistry of anthocyanin

Anthocyanin at acidic pH <2. 5 have good ground-state chemistry and they give the study of anthocyanin in aqueous solution and when it is in the hydrolysable fluid then only it gives the ground state. It usually forms the AH<sup>+</sup> flavylium cation that is pH 3, they occur the nucleophilic attack then the loss of proton give the yellow color with the hemiacetal and hemiketal form B (19). Ring-opening tautomerism can occur for ring B it forms hydroxy chalcone and Z-hydroxychalcone. Which will be isolizable thermolizable and and it is formally the state when multiple equilibrium is needed (9). Conjugate pH which raises quickly form conjugate base. Then with water, the A can observe the pH initially nucleophilic attack at 7 pH. Additional colours for the additional deprotons and losses the main colour and hence stabilized by the effective loss of pigment stabilization. This requires because of the food colour and pigment as they are much reliable on the food customer based product (20).




#### Classification of anthocyanin




Anthocyanin its glycoside and anthocyanin chemical structure is shown here

3 glucosides of the anthocyanidins are the main functional sugar group. Anthocyanidin is bifurcated sometimes sugar free anthocyanins aglycones and glycones. Literature data shows as of now 58 are reported in literature later the amount is increased more or less five hundred different anthocyanins example wise but the chemical structure is 59 moiety based they have different derivatives but the core scaffold is same shown over here. this are all literature data (14) (21) (22) (20).

Structural moieties of anthocyanin class are given below-



#### Common anthocyanin and anthocyanidin derivatives



#### Biological activity

Attractive colour is one of the most salient features of anthocyanin, like red, blue. When they are in plants the attractiveness shows cross pollination and protect and harmful effect of UV (24). Some colorant company they used to manufacture colour by these natures anthocyanin. they have a highest use in horticulture (25). Sometimes in the structure variety they show that B ring substitution glycosylation acylation with the cinnamic acid for the pigment stability. Anthocyanin pigments are very much effective and they have much effect in the radical scavenging activity for humans. Anthocyanins used as an anti oxidant and these have considerable health benefit anthocyanin based drugs nutrient are in clinical trials and they are also considered by different regulatory agencies of different companies (26).

**Bioactivity of anthocyanin pigments**

Intensified data of anthocyanins as a pigment shows that this can reduce the risk of coronary arterial disease, stroke, anti carcinogenic effect, anti Inflammatory, beneficial for improving visual acuity level and last but not the least cognitive function (27). Literature says this has a good anecdotal accounts this are also having good folklore action for the ethnic people and as a tribal medicine. For improved night vision and visual acuity the RAF force of british army they are treated with bilberries (28), Chernobyl victims feeded by Aronia melanocarpa(chokeberry) products to ameliorate effects of gamma radiation exposure. Components of red wine is highly accepted for the polyphenolic groups and due to that the antioxidant activity (5). The anthocyanin pigments are good scavenging activity and have a free radical scavenging activity towards the ROS, which have been demonstrated in vitro through such assays as oxygen radical absorbing capacity(ORAC). Various literatures show that cyanidin have much more activity than malvidin and malvidin have some less activity that pelargonidin. The effect of glycosidic substitution and acylation on the Oxygen radical absorbing capacity which is resulted in a marked increase in ORAC assay. Glycosidic substitution influenced ORAC activity, but not in a predictable manner (7). in vivo study with animals and there behavioural test are convincing. Various reviews shows that red cabbage they reduce the tumour of a rat, that are also have been due to the effect of anthocyanin (29).

This effect that is improved by dietary supplement with blueberry and strawberry extracts. In Italy, they develop one clinical study on that study they(160 mg twice daily for 1 month) exhibited improvement of diabetic retinopathy and nephropathy at the end of the trial (30).Intact content of anthocyanin glycoside found in human urine (32).

**Different biological activity of Anthocyanin**




Figure 3) Diffrent biological activity of anthocyanin.

**Biosynthetic pathway**

**Biosynthetic pathway overview**

Like other biochemical pathway anthocyanin pathway has a responsibility to produce anthocyanin pigments, the plants which give anthocyanin pigments tissue. This pathway consists a group of genes and the enzymes that are very much relevant (39). total 6 genes are core they diversified taxonomically and some other are clones by genetic mutation. The plant species Maize and snapdragon have the 2 different type of genes, literature shows that they have also cloned the 3rd one, the common morning glory, Ipomoea purpurea (40). The additional information provides the gene sequencing among the single genetic pathway (41). The genes shows different evolutionary pathway. The adaptive gene takes the pathway some other way. Regulatory gene associated with the anthocyanin pathway the adaptive gene is for the flower colour and the pigments. Sometimes structural genes also

have some capability to show function in increase vacuolar fluid and details. For the individual anthocyanin genes these 3 genes heterogeneity is seen in major angiosperms. Many times CHS type gene are in indicating without co ordinate the adjusting pathway (42) (43).

Figure mentioned anthocyanin biosynthetic pathway in angiosperms, the enzymes are CHS - chalcone synthetase CHI - Chalcone flavones isomerise, F3H flavanone isomrase, F3H flavones hydroxylase, DFR - Dihydroflavonol reductase, ANS- Anthocyanidin synthetase, and UDP3GT - UDP glucose flavonoid 3 oxy glucosyltransferase (44)




Figure 4) Biosynthetic pathway overview.

**The biosynthetic mechanism of anthocyanin pathway -**

Biosynthetic pathway of anthocyanin the well established as a network of many pathways. This pathway correlates with the flavonoid pathway also starts with chalcone synthase synthesized naringenin chalcone from four coumaryl CoA and malonyl - CoA. After that chalcone is being isomerized by naringenin. It isomerise CHI from naringenin. Naringenin is converted by the flavonoid 3'-hydroxylase or flavonoid 3, 5 hydroxylase into 2 other biflavonoids, that is dihydroquercetin and dihydromyrecetin. Respectively the next 3 pathway are same fashion. By DFR Dihydroquercetin followed by leucocyanidin by ANS then cyanidin then by biosynthetic process they have done the leucodelphinidin (45). Cyanidin pelargonidin delphinidin converts to anthocyanidin and cyaniding 3 glucosidase pelargonidin 3 glucosidase and delphinidin 3 glucosides form the anthocyanin (46). The 2 enzyme F3H and F'3'5H are diversified and determined by the B ring hydroxylation and sometimes DFR also responsible for the composition and pigmentation. This diversification varies through genus species based, altogether they all having the same biosynthetic pathway with some enzymes and all same process fields and mechanism is alters (47). But the individual steps mechanisms of action were different as oxidation reduction occurred and the coloured interaction of each level has been changed (41).

**Chemistry of anthocyanin degradation**

pH solution have a capacity to show colour in different pH. Red colour will show in pH 1 the flavylium cation. Sometimes they show blue and red colour also due to the variation of pH and the change in metal ion and the complexations. pH 2, 4 for purple and red. the quinoidal colour that is the blue predominath when ph 5 to 6, that is the carbinol pseudobase and a chalcone (3). pH value sometime 7 when the anthocyanins groups are degraded. Then the pH value is changed mainly structural pH are 4 to 6 normally seen in yellow chalcone. Flavylium is helped out when it is cation of or pH is increased. like the litmus theory it always shows the normal colouration The groups that are in ring A and B are much more effective in changing of this colour that is why when derivatization occurs it shows the same theory (55). Pelargonidin that is the most stable based on there aglycone property and monoglycosides and diglycosides are available as a

stable anthocyanin. In degradation reaction they follow the colour change basis of phenolic and aldehyde compound mainly basis of degradation reaction. The stability of anthocyanin is also the pH specific (35, 58).

#### Anthocyanin in clinical trials

Anthocyanin has different phase of action against different field of disease among them diabetes is one of them, another antioxidant, anti obesity, there are different derivatives of anthocyanin are associated among them some are good clinical agent too (60). Molecules that are under clinical trials are different type of majorly anthocyanin molecules in the field of anti obesity and anti diabetic they are in clinical trials they show the pathway in different pathway. Majorly the standardized drug are reported for the AMPK phosphorylation and the GLUT4, CRP-1, PPAR- $\alpha$  IRS-1 pathway (61). They follow sometimes AKT pathway some are WNT. Sometimes used through gene expression RBP4 and SREBP expression. mRNA based expression system is also there for inhibiting the fatty acid and the triglycerol synthesis enzyme. The downregulation always by the lipogenic factors (62).

Anthocyanin stability is depend on different buffer solutions and during storage. Different storage condition and pH 1.0,4,5 is showed still attractive colour in red and violet. Depending upon of the colour of anthocyanin , anthocyanin degradation and its storage is first order kinetics. The reaction rate( $r$ ) expressed first order and the conc based on single reactant (A) which raised to first power (72).

$$r = d[A]/dt = k[A]$$

k is the first order kinetics unit is S<sup>-1</sup>

Arranging the equations

$$d[A]/dt = -kdt$$

pH and the stability study with heat is useful for the study of degradation chemistry and other anthocyanin. Anthocyanin and anthocyanidins as a natural dye , these coloured pigment and potential colour are most stable of anthocyanin class , because they have a solubility issue. Anthocyanin have several sub class and they are potential colouring substance , yield wise in fruit also they have 50%,12%,12%,7%,7% of cyaniding, delphinidin, pelargonidin, peonidin, malvidin and petunidin , respectively. Their stability basically depend upon the pH and their light stability copigments, pigments , temperature , pH metal ions , oxygens , antioxidants. The B ring of anthocyanin moiety is mainly responsible for the stability in solution. That is why the colour changes in solution depend the anthocyanin stability. and successively the colour is depend upon the pH. Polymerization of anthocyanins due to tannins is also used to increase colour stability at lower pH. These colouring agents are much more appreciable in case of their food additives USFDA also approved anthocyanin as a colouring agents (73).

#### Root cause analysis

Anthocyanin, nature's pigment they are the normal phytoconstituent. It has a large use in the field of pigment industry , colour industry. To make food lustrous, glossy, colouring agents are sometimes used for these purpose. But for drug candidate for the best therapeutic activity it is not still been explored, some potential candidate have been through the drug discovery phases but fall down. The root cause of this is their stability, as we know the anthocyanin has the sub class of flavonoid having 3 aromatic ring and the middle ring have one ion that is positive cation. It is sometimes the flavylium ion. If we see any anthocyanin product stability we can easily see that the quinoid base have some chemical reaction, specially the oxidation and the reduction in presence of air. Quinoid base is reduced and it form the flavylium ion. then it get hydrolysed by moisture and form the hemiketal compound. The hemiketal compound then moved to the chalcone formation and chalcone can not be a anthocyanin it is a flavonoid which is basically toxic. Further addition if we take this chalcone molecule also it can also further sub divided and form the phloroglucinol aldehyde and the protocatechuic acid, means the chalcone moiety is getting reduced and form the aldehyde and carboxylic acid (74).

Here the scheme of the degradation of the hemiketal form to the flavylium ion

Beneath that we have shown the formation of aldehyde and carboxylic acid

formation. So as we can see the anthocyanin molecule has a large chance to getting reduced. So if we used it as a potential candidate and the basic structure then it may degrade the final product. In the pharmaceutical formulation study there are some class of drug they are high permeability but low solubility largely known as BCS class 2. The anthocyanin formulations are BCS class 2 drugs and that is why. This class of drugs can not be all time diluted or soluble they have some issue of solubility also. That is why by overcoming all this obstacles it is very hard to develop anthocyanin as a potential drug candidate. But technology develops , a large class of drugs are now administered through nano formulation and nano crystals. it can help in stability and there is a possibility of potential drug candidate (75).

An acylated cyanidingly coside in the red - purple of *Xanthoxanthin* cvminipurple

#### Future perspectives

Metabolic property all time not supplied by our body some times it is needed based on the physiological condition , sometimes they are isomers and they have same metabolic property. The anthocyanin and anthocyanidin drug development story is such kind of chemical instability problem which are against of some natural properties. It has also some biphasic conditions for acceptable period and it also helps for chemoprevention anti carcinogenic robust , and also helps modify of studies. These are the sort of anti obesity drugs also a clinical candidate of anthocyanins, is clearly superior parts of anti obesity drugs. Presently pre clinical and clinical differences and the questions conclusively if the mixtures developed the appropriate strategy have show anthocyanin as a good pharmacokinetic agent. Though some stability problem arises in many case of compounds don't have proper stability and the mixture contains some degradants. Presently the anthocyanin products that are in market are well established , researchers are doing study for there components profile of individual mixture of components. Study also shown that the mixture can also the thing which contains fruits and many components may contribute to the chemopreventive efficacy.

#### REFERENCES

- Kendrick A. Coloring Aqueous Food Types. Handbook on Natural Pigments in Food and Beverages: Industrial Applications for Improving Food Color. Elsevier Ltd; 2016. 163-177 p.
- Wrolstad RE. Natural Food colours, Science and technology. J Food Sci. Elsevier Ltd; 2000;14:C419-25.
- Union E, Physiological C, Genetic B, Further R. Anthocyanin Occurrence of anthocyanins. Hindawi publishing house; 2014; 11-36pg
- Konczak I, Zhang W. Anthocyanins – More Than Nature's Colour. 2004;5:239-40.
- Farzaei MH, El-Senduny FF, Momtaz S, Parvizi F, Iranpanah A, Tewari D, et al. An update on dietary consideration in inflammatory bowel disease: anthocyanins and more. Expert Rev Gastroenterol Hepatol, Hiral publisher ;123-632
- de Pascual-Teresa S, Moreno DA, García-Viguera C. Flavanols and anthocyanins in cardiovascular health: A review of current evidence. Int J Mol Sci. 2010;11 (4):1679-703.
- Gertrude BY, Robinson M, Robinson R. Cxxxii. a survey of anthocyanins. i. Biochemistry. 1931;1687-705.
- He J, Giusti MM. Anthocyanins: Natural Colorants with Health-Promoting Properties. Annu Rev Food Sci Technol. 2010;1 (1):163-87.
- Strack d, Wray v. Anthocyanins. 1989;325-56.
- Bullerjahn GS, Cable JR, Castellano FN, Glusac K, Lu HP, Ogawa MY, et al. Editor. 2006;19 (3).
- T. Goto TK. - 61 1.1. Agric Chem. 1991;17-33.
- Brouillard R, Dubois JE. Mechanism of the Structural Transformations of Anthocyanins in Acidic Media. J Am Chem Soc. 1977;99 (5):1359-64.
- García-Viguera C, Bridle P. Influence of structure on colour stability of anthocyanins and flavylium salts with ascorbic acid. Food Chem. 1999;64 (1):21-6.
- Silva VO, Freitas AA, Maçanita AL, Quina, Frank H. Silva, V. O. et al. (2016) "Chemistry and photochemistry of natural plant pigments:

- the anthocyanins”, *Journal of Physical Organic Chemistry*, 29 (11) pp. 594–599.
15. Smeriglio A, Barreca D, Bellocco E, Trombetta D. Chemistry, Pharmacology and Health Benefits of Anthocyanins. *Phyther Res*. 2016; (April):1265–86.
  16. Amr A, Al-Tamimi E. Stability of the crude extracts of *Ranunculus asiaticus* anthocyanins and their use as food colourants. *Int J Food Sci Technol*. 2007;42 (8):985–91.
  17. Boots AW, Haenen GRMM, Bast A. Health effects of quercetin: From antioxidant to nutraceutical. *Eur J Pharmacol*. 2008;585 (2–3):325–37.
  18. Chumbhale DS, Chaudhari SR, Upasani CD. Available online at www.ijpcbs.com Research Article Standardization of marketed kumari asava-a polyherbal ayurvedic formulation. 2014;4 (3):681–5.
  19. F. DelgadoVargas ARJ& OPL. Natural Pigments : Carotenoids , Anthocyanins , and Natural Pigments : Carotenoids , Anthocyanins , and Betalains – Characteristics , Biosynthesis , Processing , and Stability. Vol. 40:3, *Critical Reviews in Food Science and Nutrition*. 2000. 173-289 p.
  20. Iacobucci GA, Sweeny JG. The chemistry of anthocyanins, anthocyanidins and related flavylium salts. *Tetrahedron*. 1983;39 (19):3005–38.
  21. Prior RL, Wu X. Anthocyanins : Structural characteristics that result in unique metabolic patterns and biological activities \*. 2006;40 (October):1014–28.
  22. Castañeda-ovando A, Pacheco-hernández MDL, Páez-hernández ME, Rodríguez JA, Galán-vidal CA. Chemical studies of anthocyanins : A review. *Food Chem*, 2009;113 (4):859–71.
  23. Wrolstad RE. Anthocyanins by UV-Visible Spectroscopy. 2001;1–13.
  24. Al-Awwadi N, Azay J, Pouchere P, Cassanas G, Krosniak M, Auger C, et al. Antidiabetic Activity of Red Wine Polyphenolic Extract, Ethanol, or Both in Streptozotocin-Treated Rats. *J Agric Food Chem*. 2004;52 (4):1008–16.
  25. Wojdyło A, Nowicka P, Carbonell-Barrachina ÁA, Hernández F. Phenolic compounds, antioxidant and antidiabetic activity of different cultivars of *Ficus carica* L. fruits. *J Funct Foods*. 2016;25:421–32.
  26. Vareed SK, Reddy MK, Schutzki RE, Nair MG. Anthocyanins in *Cornus alternifolia*, *Cornus controversa*, *Cornus kousa* and *Cornus florida* fruits with health benefits. *Life Sci*. 2006;78 (7):777–84.
  27. Martineau LC, Couture A, Spoor D, Benhaddou-Andaloussi A, Harris C, Meddah B, et al. Anti-diabetic properties of the Canadian lowbush blueberry *Vaccinium angustifolium* Ait. *Phytomedicine*. 2006;13 (9–10):612–23.
  28. Gould KS. Nature ’ s Swiss Army Knife : The Diverse Protective Roles of Anthocyanins in Leaves. 2004;5:314–20.
  29. Shih PH, Wu CH, Yeh CT, Yen GC. Protective effects of anthocyanins against amyloid  $\beta$ -peptide-induced damage in neuro-2A Cells. *J Agric Food Chem*. 2011;59 (5):1683–9.
  30. Wang H, Nair MG, Strasburg GM, Chang Y, Booren AM, Gray JJ, et al. P & A - 137 Antioxidant and Antiinflammatory Activities of Anthocyanins and Their. *Reactions*. 1999; (517):294–6.
  31. Grace MH, Ribnicky DM, Kuhn P, Poulev A, Logendra S, Yousef GG, et al. Hypoglycemic activity of a novel anthocyanin-rich formulation from lowbush blueberry, *Vaccinium angustifolium* Aiton. *Phytomedicine*. 2009;16 (5):406–15.
  32. Wang LS, Stoner GD. Anthocyanins and their role in cancer prevention. *Cancer Lett*. 2008;269 (2):281–90.
  33. Nizamutdinova IT, Jin YC, Chung J, Shin SC, Lee SJ, Seo HG, et al. The anti-diabetic effect of anthocyanins in streptozotocin-induced diabetic rats through glucose transporter 4 regulation and prevention of insulin resistance and pancreatic apoptosis. *Mol Nutr Food Res*. 2009;53 (11):1419–29.
  34. Feild TS, Lee DW, Holbrook NM. Why Leaves Turn Red in Autumn. The Role of Anthocyanins in Senescing Leaves of Red-Osier Dogwood. *Plant Physiol*. 2001;127 (2):566–74.
  35. Coman C, Rugină OD, Socaciu C. Plants and natural compounds with antidiabetic action. *Not Bot Horti Agrobot Cluj-Napoca*. 2012;40 (1):314–25.
  36. Kameswara Rao B, Renuka Sudarshan P, Rajasekhar MD, Nagaraju N, Appa Rao C. Antidiabetic activity of *Terminalia pallida* fruit in alloxan induced diabetic rats. *J Ethnopharmacol*. 2003;85 (1):169–72.
  37. Lopez-Martinez LX, Oliart-Ros RM, Valerio-Alfaro G, Lee CH, Parkin KL, Garcia HS. Antioxidant activity, phenolic compounds and anthocyanins content of eighteen strains of Mexican maize. *LWT - Food Sci Technol*. 2009;42 (6):1187–92.
  38. You Q, Chen F, Wang X, Jiang Y, Lin S. Anti-diabetic activities of phenolic compounds in muscadine against  $\alpha$ -glucosidase and pancreatic lipase. *LWT - Food Sci Technol*. 2012;46 (1):164–8.
  39. Yoder JI, Belzile F, Tong Y, Goldsbrough A. Visual markers for tomato derived from the anthocyanin biosynthetic pathway. *Euphytica*. 1994;79 (3):163–7.
  40. Katsumoto Y, Fukuchi-Mizutani M, Fukui Y, Brugliera F, Holton TA, Karan M, et al. Engineering of the rose flavonoid biosynthetic pathway successfully generated blue-hued flowers accumulating delphinidin. *Plant Cell Physiol*. 2007;48 (11):1589–600.
  41. Winkel-shirleyB. Winkel-Shirley2001-flavonoid-color-biosynthesis. pdf2001;126 (June):485–93.
  42. Evans O. “Ein botschafter des vergessenen”: de bruyn, the chronicler. *Ger Monit*. 2012;75 (8):185–97.
  43. He F, Mu L, Yan GL, Liang NN, Pan QH, Wang J, et al. Biosynthesis of anthocyanins and their regulation in colored grapes. *Molecules*. 2010;15 (12):9057–91.
  44. Tanaka Y, Sasaki N, Ohmiya A. Biosynthesis of plant pigments: Anthocyanins, betalains and carotenoids. *Plant J*. 2008;54 (4):733–49.
  45. Goff SA, Klein TM, Roth BA, Fromm ME, Cone KC, Radicella JP, et al. Transactivation of anthocyanin biosynthetic genes following transfer of B regulatory genes into maize tissues. *EMBO J*. 1990;9 (8):2517–22.
  46. Solfanelli C, Poggi A, Loreti E, Alpi A, Perata P. 1’a004Mmcr. 2006;1:637–746.
  47. Loreti E, Povero G, Novi G, Solfanelli C, Alpi A, Perata P. Gibberellins, jasmonate and abscisic acid modulate the sucrose-induced expression of anthocyanin biosynthetic genes in *Arabidopsis*. *New Phytol*. 2008;179 (4):1004–16.
  48. Sui X, Zhang Y, Zhou W. In vitro and in silico studies of the inhibition activity of anthocyanins against porcine pancreatic  $\beta$ -amylase. *J Funct Foods*. 2016;21:50–7.
  49. Liu Y, Tikunov Y, Schouten RE, Marcelis LFM, Visser RGF, Bovy A. Anthocyanin Biosynthesis and Degradation Mechanisms in Solanaceous Vegetables: A Review. *Front Chem*. 2018;6.
  50. Ben-Yehudah G, Korchinsky R, Redel G, Ovadya R, Oren-Shamir M, Cohen Y. Colour accumulation patterns and the anthocyanin biosynthetic pathway in “Red Delicious” apple variants. *J Hort Sci Biotechnol*. 2005;80 (2):187–92.
  51. Moss GP. Flavonoid biosynthesis. Web version *Enzym Nomencl*. 2001;2007 (April 30, 2007):Web version edited from *Enzyme Nomenclature* 1992:1145-1896.
  52. Khoo HE, Azlan A, Tang ST, Lim SM. Anthocyanidins and anthocyanins: colored pigments as food, pharmaceutical ingredients, and the potential health benefits. *Food Nutr Res* 1998:456-678.
  53. Holt RR, Lazarus SA, Sullards MC, Zhu QY, Schramm DD, Hammerstone JF, et al. Plasma After the Consumption of a Flavanol-Rich Cocoa. *Am J Clin Nutr*. 2002; (February):798–804.
  54. Hosseinian FS, Li W, Beta T. Measurement of anthocyanins and other phytochemicals in purple wheat. *Food Chem*. 2008;109 (4):916–24.
  55. Bassa la, francis fj. Stability of Anthocyanins from Sweet Potatoes in a Model Beverage. *J Food Sci*. 1987;52 (6):1753–4555.
  56. Sano a, yamakoshi j, tokutake s, tobe k, kubota y, kikuchi m. Procyandin B1 Is Detected in Human Serum after Intake of Proanthocyanidin-rich Grape Seed Extract. *Biosci Biotechnol Biochem*. 2003;67 (5):1140–3789.
  57. You Q, Chen F, Wang X, Luo PG, Jiang Y. Inhibitory effects of muscadine anthocyanins on  $\alpha$ -glucosidase and pancreatic lipase activities. *J Agric Food Chem*. 2011;59 (17):9506–11.
  58. Newman DJ, Cragg GM. Natural Products as Sources of New Drugs from 1981 to 2014. *J Nat Prod*. 2016;79 (3):629–61.

59. Rausher MD, Miller RE, Tiffin P. Patterns of evolutionary rate variation among genes of the anthocyanin biosynthetic pathway. *Mol Biol Evol.* 1999;16 (2):266-74.
60. Azzini E, Giacometti J, Russo GL. Antiobesity Effects of Anthocyanins in Preclinical and Clinical Studies. *Oxid Med Cell Longev.* 2017;2017 (ii).
61. Guenther BD, Christensen CR, Upatnieks J. Coherent Optical Processing: Another Approach. *IEEE J Quantum Electron.* 1979;15 (12):1348-62.
62. Núñez-Sánchez MA, González-Sarriás A, Romo-Vaquero M, García-Villalba R, Selma M V, Tomás-Barberán FA, et al. Dietary phenolics against colorectal cancer-From promising preclinical results to poor translation into clinical trials: Pitfalls and future needs. *Mol Nutr Food Res.* 2015;59 (7):1274-91.
63. Lila MA. Anthocyanins and Human Health : An In Vitro Investigative Approach. 2007;2004 (5):1-13.
64. Pojer E, Mattivi F, Johnson D, Stockley CS. The case for anthocyanin consumption to promote human health: A review. *Compr Rev Food Sci Food Saf.* 2013;12 (5):483-508.
65. Feng R, Ni HM, Wang SY, Tourkova IL, Shurin MR, Harada H, et al. Cyanidin-3-rutinoside, a natural polyphenol antioxidant, selectively kills leukemic cells by induction of oxidative stress. *J Biol Chem.* 2007;282 (18):13468-76.
66. Belwal T, Nabavi SF, Nabavi SM, Habtemariam S. Dietary anthocyanins and insulin resistance: When food becomes a medicine. *Nutrients.* 2017;9 (10).
67. Pasqualone A, Bianco AM, Paradiso VM. Production trials to improve the nutritional quality of biscuits and to enrich them with natural anthocyanins. *CYTA - J Food.* 2013;11 (4):301-8.
68. C. Patent Application Publication( 10 ) Pub. No.: US 2004 / 0109905 A1. 2004;1 (19).
69. Mazza G, Brouillard R. Recent developments in the stabilization of anthocyanins in food products. *Food Chem.* 1987;25 (3):207-25.
70. Lizasoain A, Tort LF, Garc'ia M, Gomez MM, Leite JP, Miagostovich MP, et al. No. *J Appl Microbiol.* 2015;119 (3):859
71. García-Viguera C, Bridle P. Influence of structure on colour stability of anthocyanins and flavylum salts with ascorbic acid. *Food Chem.* 1999;64 (1):21-6.
72. Torskangerpoll K, Andersen ØM. Colour stability of anthocyanins in aqueous solutions at various pH values. *Food Chem.* 2005;89 (3):427-40.
73. Oancea S, Drăghici O. PH and thermal stability of anthocyanin-based optimised extracts of romanian red onion cultivars. *Czech J Food Sci.* 2013;31 (3):283-91.
74. Fischer UA, Carle R, Kammerer DR. Thermal stability of anthocyanins and colourless phenolics in pomegranate(*Punica granatum L.*) juices and model solutions. *Food Chem.* 2013;138 (2-3):1800-9.
75. Thomasset S, Teller N, Cai H, Marko D, Berry DP, Steward WP, et al. Do anthocyanins and anthocyanidins, cancer chemopreventive pigments in the diet, merit development as potential drugs; *Cancer Chemother Pharmacol.* 2009;64 (1):201-11
76. Asen, S., Stuart, N.W., and Siegelman, H.W. 1959. Effect of various concentrations of nitrogen, phosphorus and potassium on sepal color of *Hydrangea macrophylla*. *Am. Soc. Hort. Sci.* 73:495-502.
77. Bridle, P. and Timberlake, C.F. 1996. Anthocyanins as natural food colors-selected aspects. *Food Chem.* 58:103-109.
78. Brouillard, R. and El Hache Chahine, J.M. 1980. Chemistry of anthocyanin pigments. 6. Kinetic and thermodynamic study of hydrogen sulfite addition to cyanin. Formation of a highly stable Meisenheimer-type adduct derived from a 2- phenylbenzopyrylium salt. *J. Am. Chem. Soc.* 102:5375-5378.
79. Dangles, O., Saito, N., and Brouillard, R. 1993. Anthocyanin intramolecular copigment effect. *Phytochemistry* 34:119-124.
80. Figueiredo, P., Elhabiri, M., Saito, N., and Brouillard, R. 1996. Anthocyanin intramolecular interactions. A new mathematical approach to account for the remarkable colorant properties of the pigments extracted from *Matthiola incana*. *J. Am. Chem. Soc.* 118:4788-4793
81. Francis, F.J. 1982. Analysis of anthocyanins. In *Anthocyanins as Food Colors*(P. Markakis, ed.) New Yorks academic press;pp. 182-205.
82. Francis, F.J. 1989. Food colorants: Anthocyanins. *Crit. Rev. Food Sci. Nutr.* 28:273-314.
83. Fuleki, T. and Francis, F.J. 1968a. Quantitative methods for anthocyanins. 1. Extraction and de- termination of total anthocyanin in cranberries. *J. Food Sci.* 33:72-78.
84. Fuleki, T. and Francis, F.J. 1968b. Quantitative methods for anthocyanins. 2. Determination of total anthocyanin and degradation index for cranberry juice. *J. Food Sci.* 33:78-82.
85. Giusti, M.M., Rodriguez-Saona, L.E., Baggett, J.R., Reed, G.L., Durst, R.W., and Wrolstad, R.E. 1998. Anthocyanin pigment composition of red radish cultivars as potential food colorants. *J. Food Sci.* 63:219-224.
86. Giusti, M.M., Rodriguez-Saona, L.E., and Wrolstad, R.E. 1999. Spectral characteristics, molar absorptivity and color of pelargonidin derivatives. *J. Agric. Food Chem.* 47:4631-4637
87. Harborne, J.B. 1967. *Comparative Biochemistry of the Flavonoids.* Academic Press, London:456-895
88. Heredia, F.J., Francia-Aricha, E.M., Rivas-Gonzalo, J. C., Vicario, I.M., and Santos-Buelga, C. 1998. Chromatic characterization of anthocyanins from red grapes. I. pH effect. *Food Chem.* 63:491-498.
89. Hong, V. and Wrolstad, R.E. 1990. Use of HPLC separation/ photodiode array detection for characterization of anthocyanins. *J. Agric. Food Chem.* 38:708-715.
90. Hrazdina, G., Iredale, H., and Mattick, L.R. 1977. Anthocyanin composition of Brassica oleracea cv. Red Danish. *Phytochemistry* 16:297-301
91. Jackman, R.L. and Smith, J.L. 1996. Anthocyanins and betalains. In *Natural Food Colorants*, 2nd ed.(G.A.F.Hendry and J.D. Houghton, eds.) Chpt.Blackie & Son, Glasgow, Scotland.
92. Jackman, R.L., Yada, R.Y., and Tung, M.A. 1987. A review: Separation and chemical properties of anthocyanins used for their qualitative and quantitative analysis. *J. Food Biochem.* 11:279-308.
93. Jorgensen, E.C. and Geissman, T.A. 1955. The chemistry of flower pigmentation in *Antirrhinum majus* color genotypes. III. Relative anthocyanin and aurone concentrations. *Biochem. Biophys.* 55:389-402.
94. Jurd, L. and Asen, S. 1966. The formation of metal and "co-pigment" complexes of cyanidin 3-glu- coside. *Phytochemistry* 5:1263-1271.
95. Kamei, H., Kojima, T., Hasegawa, M., Koide, T., Umeda, T., Yukawa, T., and Terabe, K. 1995. Suppression of tumor cell growth by anthocyanins in vitro. *Cancer Invest.* 13:590-594
96. Karaivanova, M., Drenska, D., and Ovcharov, R. 1990. A modification of the toxic effects of platinum complexes with anthocyanins. *Eksp. Med. Morfol.* 29:19-24.
97. Kraemer-Schafhalter, A., Fuchs, H., Strigl, A., Silhar, S., Kovac, M., and Pfannhauser, W. 1996. Process consideration for anthocyanin extraction from Black Chokeberry(*Aronia meloncarpa* ELL). In *Proceedings of the Second International Symposium on Natural Colorants*, INF/COL II(P.C. Hereld, ed.), pp.153-160.
98. Mazza, G. and Miniati, E. 1993. Introduction. In *Anthocyanins in fruits, vegetables, and grains.*(G. Mazza and E. Miniati, eds). CRC Press, Boca Raton, Fla.45thEdn,145-134.
99. McClure, J.W. 1967. Photocontrol of *Spirodela intermedia* flavonoids. *Plant Phys.* 43:193-200.
100. Metivier, R.P., Francis, F.J., and Clydesdale, F.M. 1980. Solvent extraction of anthocyanins from wine pomace. *J. Food Sci.* 45:1099-1100
101. Niketic-Aleksic, G. and Hrazdina, G. 1972. Quantitative analysis of the anthocyanin content in grape juices and wines. *Lebensm. Wiss. U. Tech- nol.* 5:163-165.
102. Ribereau-Gayon, P. 1959. *Recherches sur les anthocyanes des vegetaux. Application au genre Vitis.* Doctoral dissertation, p.118.
103. Sakamura, S. and Francis, F.J. 1961. The anthocyanins of the American cranberry. *J. Food Sci.* 26:318-321.
104. Schou, S.A. 1927. Light absorption of several anthocyanins. *Helv. Chim. Acta.* 10:907-915.

105. Siegelman, H.W. and Hendricks, S.B., 1958. Photo-control of alcohol, aldehyde and anthocyanin production in apple skin. *Plant Phys.* 33:409-413.
106. Somers, T.C. 1966. Grape phenolics: the anthocyanins of *Vitis vinifera*, var. Shiraz. *J. Sci. Food Agric.* 17:215-219.
107. Somers, T.C. and Evans, M.E. 1974. Wine quality: Correlations with colour density and anthocyanin equilibria in a group of young red wines. *J. Sci. Food Agric.* 25:1369-1379.
108. Strack, D. and Wray, V. 1989. Anthocyanins. In *Methods in Plant Biochemistry*, Vol. I, Plant Phenolics (P.M. Dey and J.B. Harborne, eds.). Academic Press, San Diego;12:1335-1345.
109. Strack, D. and Wray, V. 1994. The anthocyanins. In *The Flavonoids: Advances in Research Since 1986* (J.B. Harborne, ed.). Chapman and Hall.34;156-589.
110. Swain, T. 1965. Analytical methods for flavonoids. In *Chemistry and Biochemistry of Plant Pigments* (T.W. Goodwin, ed.). Academic Press, London.45:36-58.
111. Swain, T. and Hillis, W.E. 1959. The phenolic constituents of *Prunus domestica*. I. The quantitative analysis of phenolic constituents. *J. Sci. Food Agric.* 10:63-68.
112. Takamura, H. and Yamagami, A. 1994. Antioxidative activity of monoacylated anthocyanins isolated from Muscat Bailey A grape. *J. Agric. Food Chem.* 42:1612-1615.
113. Timberlake, C.F. 1980. Anthocyanins [related to beverages]—occurrence, extraction and chemistry [coloring material]. *Food Chem.* 5:69-80
114. Timberlake, C.F. and Henry, B.S., 1988. Anthocyanins as natural food colorants. *Prog. Clin. Biol. Res.* 280:107-121.
115. Wang, H., Cao, G., and Prior, R.L., 1997. Oxygen radical absorbing capacity of anthocyanins. *J. Agric. Food Chem.* 45:304-309
116. Wrolstad, R.E., Putnam, T.P., and Varseveld, G.W. 1970. Color quality of frozen strawberries: Effect of anthocyanin, pH, total acidity and ascorbic acid variability. *J. Food Sci.* 35:448-452.
117. Wrolstad, R.E., Hong, V., Boyles, M.J., and Durst, R.W. 1995. Use of anthocyanin pigment analysis for detecting adulteration in fruit juices. In *Methods to Detect Adulteration in Fruit Juice and Beverages*, 23:449-344.
118. Patent details Reference
119. Kim Hyeong Jeong Shin Sung Chul Chong Chong Il, 2009. Composing controlling diabetic condition. KR101190624B1.25. 05.2009.
120. Mitsunori Ono, 2006. Composition for treating diabetes or obesity, US20060188590A1.18.01.2006.
121. David Ribnicky, 2009. Berry preparation for treating diabetes and metabolic syndrome. US20090176718A1.17.01.2009.
122. Jang Bong Keun Han Sang Jin, 2004. Composition containing blackchokeberry extracts for diabetes treatment. KR20100036144A07.04.2004
123. Mark Andrew Paulik, 2014. Composition and methods for treating diabetes and obesity. US20140349923A1.09.03.2014.
124. Mike Farber, 2008. Anthocyanins, shown to have an effect on endothelial nitric oxide. US20080063689A1.03.06.2008.
125. Barney W. Hilto, 2009. Processed anthocyanin pigment extracts. US4320009A04.04.2009.
126. Enno Krebbers, Use of anthocyanin gene in male sterile plants. US5880331A.14.12.2008.
127. Wayne H. Crosby, 2004. Stabilized anthocyanin as food colorant, United States patent. US4481226A.30.10.2004.
128. Metty S. K. Langston, 2005. Anthocyanin colorant from grape pomace, US4500556A.05.05.1995
129. Bruce K. Wallin, 2001. Process for purifying anthocyanin colours. US4211577A.23.02.2001
130. Thomas Philip, 2005. Recovery of anthocyanin from plant source US3963700A.12.12.2005.
131. Isao Yokoyama, 2009. Process of extracting natural colours from anthocyanin as natural products. US4302200A.01.05.2009.
132. Anil J. Shrikhand, 2010. Extraction and intensification of anthocyanin from grape pomace. US4452822A.10.10.2010.
133. Debasis Bagchi, 2004. Method and composition of anthocyanin-rich berry extracts that prevents or inhibits angiogenesis and helicobacter pylori and acts as a powerful antioxidant that provides various health benefits. US20040109905A1.09.01.2004.
134. Rod Lenoble, 2017. Pigment composition containing anthocyanins stabilized by plant extracts US5908650A.08.09.2014.
135. Lee Kang-soo Lee Hyun-hyun, 2009. Functional beverage of Omija with stable anthocyanin, Korean patent file. KR101139634B1.14.12.2009.
136. Muraleedharan Nair, 2006. Insulin secretion by anthocyanin and anthocyanidins. US20060025353A1.02.06.2006.
137. Zheng Lejian, 2006. Divaricate velvet plant medicinal preparation for treating diabetes. CN101147749A.24.06.2015.
138. Leenaning Zhopang, Method of preparing pumpkin seeds containing rich anthocyanin. CN102845788A.21.12.2006.
139. Kim Kada khyuch, 2015. Composition for preventing or treating diabetes comprising black corn plant extracts, health food comprising the same, and method for preparing the same. KR20150125339A.05.01.2015.
140. Manging jon hyun pangong. Composition for improving diabetes and bloodstream and method for preparing the same. KR101813015B1.13.12.2015.
141. Jie Luo, 2013. Method of change the gene expression of solanaceous plant for diabetes treatment, US8802925B2.15.11.1995.
142. Mark L. Heiman, 2015. Method to treat diabetes utilizing a gastrointestinal microbiome modulating composition, US20150118330A1.11.08.2015.
143. Hym zen jon. 2015. Composition for treating diabetes as well as preparation method and application of composition CN105341760A.12.11.2015.
144. Ilya Raskin, 2007. Red lettuces with increased anthocyanins, polyphenols, and oxygen radical absorption capacity. US9480235B1.14.11.2007
145. yuu tong ming hyam. 2017. Composition for improving diabetes and bloodstream and method for preparing the same, KR20170043166A.17.12.2017.
146. Chung chang hung. 2012. Orally disintegrating tablet for type 2 DM. CN106265566A10.12.2012.
147. Mark Pedersen, 2006. Metabolic capacity enhancing compositions and methods for use in a mammal, US20060024385A1.14.11.2006.
148. Stanley Kubow, 2012. Methods of treating obesity in type-2 DM, US9446063B2.17.10.2012.
149. Masayuki Yoshikawa, 2007. Novel inhibitor of the formation of advanced glycation end product and aldose reductase inhibitor, US20070060533A1.05.06.2007.
150. Guntang jah, kim jon, 2009. Sea cucumber anthocyanin flour and preparation method of vermicelli, CN102813158A.01.03.2009.
151. Tae Hyun Choi, 2008. Pharmaceutical composition for wound healing containing anthocyanin extracted from the black soybean seed coat. WO2008126980A1.06.12.2008.
152. John Doherty, 2012. Compositions and methods for promoting weight loss and increasing energy. US20120034323A1.03.04.2012.