

Commentary on Beam Computed Tomography Studies in Sphenoid Sinus

Natalisa Hvizdosova*

Hvizdosova N. Commentary on Beam Computed Tomography Studies in Sphenoid Sinus. *Int J Anat Var.* 2021;15(5):177-177.

ABSTRACT

The aim of this Commentary is to define cone beam computed tomography (CBCT) characteristics of arrested pneumatization of sphenoid sinus in an effort to help differentiate it from invasive or lytic skull base lesions. Two

cases are presented with incidental findings. Both studies, acquired for other diagnostic purposes, demonstrated unique osseous patterns that were eventually deemed to be anatomic variations in the absence of clinical signs and symptoms although the pattern of bone loss and remodeling was diagnosed as pneumatization of the sphenoid sinus by a panel of medical and maxillofacial radiologists following contrasted advanced imaging.

Key Words: *Phenoid sinus; Development; Cone-beam computed tomography*

INTRODUCTION

Since the appearance of relatively lucent lesions mimic osteolysis in the skull base captured on such studies and trigger additional investigations, it is imperative that benign conditions noted in such studies be considered in the differential diagnoses prior to further referral and advanced imaging [1,2]. This report presents two cases in patients reporting for orthodontic or orthognathic surgery and dental implant placement procedure. The aim of the report is to investigate owing to extensive effacement of the sphenoid bone architecture. This report presents two cases of arrested pneumatization of the sphenoid sinus as seen on large field-of-view (FOV) cone beam computed tomography studies (CBCT) acquired for orthodontic/maxillofacial surgical diagnostic tasks and dental implant placement procedure. The aim of the report is to define CBCT characteristics of arrested pneumatization of sphenoid sinus in an effort to help differentiate it from aggressive, invasive skull base lesions [3]. CBCT is suboptimal for interpretation of soft tissue entities and spatial resolution is significantly higher than that of multi-detector computed tomography (MDCT). Spatial resolution is better than MDCT by an order of magnitude but the modality is best suited for imaging osseous and dental tissues. Image acquisition is completed in 20 seconds, with the total radiation dose being a fraction of that associated with MDCT [4].

There were central heterogeneous, high-attenuation entities with signal intensities approximating those of osseous tissue. Possibility of involvement of vidian canal and foramen rotundum could not be ruled out. Based upon the radiographic presentation, an impression of arachnoid granulations was made [4]. There was also evidence of bowing and thinning of the lingual cortical plate at the left mandibular ramus superior to the lingula of the mandibular foramen.

Diagnosing arrested pneumatization is important in order to distinguish this entity from other skull base abnormalities like fibrous dysplasia, arachnoid granulations, and invasive lesions such as acoustic neuroma, glioma, and pituitary tumor such as an adenoma, giant cell tumor, chondro sarcoma, metastatic lesions, meningioma, or chordoma [5].

ACKNOWLEDGEMENT: None.

CONFLICTS OF INTEREST: None.

REFERENCES

1. Zeynep G, Elif B. Evaluation of sphenoid sinus variations by using cone beam computed tomography in patients with cleft lip/palate. *Oral Radiol.* 2020; 36(3):248-253.
2. José LT, Félix M. Sphenoid Sinus: Pneumatization and Septation Patterns in a Hispanic Population. *ORL J Otorhinolaryngol Relat Spec.* 2021; 83(5):362-371.
3. Beryl SO, Patrick SI. Variant neurovascular relations of the sphenoid sinus in adult Nigerians. *Niger Postgrad Med J.* 2021; 28(2):112-116.
4. Yujia Y, Fang G, Jiayu L, et al. Anatomical Variants, Pneumatization Classification, and Volumetric Studies of the Sphenoid Sinus with High-Resolution Computed Tomography. *J Craniofac Surg.* 2021; 1; 32(7):2542-2545.
5. Ziya Ş, Nuray BM. Relation between optic and carotid canals with sphenoid sinus in patients with communicant hydrocephalus: a computed tomography evaluation study. *Acta Radiol.* 2020; 61(8):1064-1071.

Department of Anatomy, Faculty of Medicine, Pavol Jozef Safarik University in Kosice, Slovak Republic

Correspondence: Natalisa Hvizdosova, Department of Anatomy, Faculty of Medicine, Pavol Jozef Safarik University in Kosice, Slovak Republic. E-mail: natalisa.hvizdosova@upjs.sk

Received: 29-Apr-2022, Manuscript No: ijav-22-4945, Editor assigned: 2-May-2022, PreQC No: ijav-22-4945 (PQ), Reviewed: 16-May-2022, QC No: ijav-22-4945, Revised: 20-May-2022, Manuscript No: ijav-22-4945 (R) Published: 27-May-2022, DOI: 10.37532/1308-4038.15(5).199



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact reprints@pulsus.com