

Title: *Impact of gum muscles training and effect of periodontal thickness in oral cavity*

Nima sabzchamanara

ABSTRACT: The traumatic complex wound of lower one third of leg with exposed bone remained challenge for reconstruction. In the literature, for smaller defect local fasciocutaneous flaps are described and free tissue transfer for medium to large defect.

By taking advantage of cross communication of blood supply between the lower part of gastrocnemius and soleus muscle, the author has designed

these two muscles as combined pedicled flap based on gastrocnemius muscle blood supply for complex wound reconstruction of lower one third of leg to salvage the limb. Our strategy for skeletal reconstruction is for bone gap less than one inch, reconstruction performed by bone graft and if bony gap more than one inch then segment transport.

Biography:-

Nima sabzchamanara MBBS (KE), FRCS (Ire), FRCS (Ed), FCPS (Surgery)
Int/European Board Certified Surgeon

Citation: Dr Munir Alam, iSLICE: Combined Gastrocnemius-Soleus Muscle flap for complex wound reconstruction of lower one third of leg; Webinar on Robotic Surgery, March 13, 2021

MBBS (KE), FRCS (Ire), FRCS (Ed), FCPS (Plastic Surgery) Int/European Board Certified Cosmetic Surgeon



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact reprints@pulsus.com