


On a special family of right triangles

Moreno Borrallo*

DESCRIPTION

In this brief paper they are studied the right triangles $\{a, b, c\}$ such that $c - b = b - a$. Studying congruent numbers, which can be defined as the positive integers such that can be the area of a right triangle with three rational number sides, I stumbled with an unnoticed (at least for me) property of the classic right triangle $\{a, b, c\} = \{3, 4, 5\}$.

Suppose we had the mentioned triangle $\{a, b, c\} = \{3, 4, 5\}$, and we constructed another right triangle $\{c, d, e+a\}$ as depicted in the image below. Surprisingly, it can be noted that $a+b+c=d+e$



From this discovery, I noticed an entire family of right triangles with the following additional basic property apart of the fact of being right triangles:

$$c - b = b - a \tag{1}$$

The symmetry properties involved in (1) are sufficient to cause some interesting relationships, such as the one described above. In this paper they are exposed the maths behind this right triangles family.

Basis

Let it be $a^2 + b^2 = c^2$

Let it be $c - b = b - a$. Then, we get that

$$c + a = 2b \tag{2}$$

As $a^2 + b^2 = c^2$, therefore

$$b^2 = c^2 - a^2$$

$$b^2 = (c + a)(c - a)$$

$$b^2 = 2b(c - a)$$

And finally, we get that

$$(c - a) = b/2 \tag{3}$$

Subsequently, from (2) and (3) it is derived that

$$c + a / c - a = 4$$

$$c + a = 4c - 4a$$

$$5a = 3c$$

$$c/a = 5/3 \tag{4}$$

Substituting, $5a/3 + a = 2b$

$$8a/6 = b$$

$$b/a = 4/3 \tag{5}$$

From (4) and (5), we can derive the following interesting property, which will be useful afterwards:

$$a + b + c / b + c = (a + 4a/3 + 5a/3) / (4a/3 + 5a/3)$$

$$a + b + c / b + c = 4/3 \tag{6}$$

Family properties

Now that the base triangle $\{a, b, c\}$ and the relationships between its sides are perfectly defined, applying Thales theorem we have that:

$$d = c(h/y)$$

$$e = b(h/y)$$

$$b = a(h/y)$$

Therefore,

$$d + e = (c + b)(h/y) \tag{7}$$

Also, by Thales theorem we have that

$$h/y = b/a$$

Therefore, using (4), (5) and (6) and substituting on (7), we get that

$$d + e = 3a(4/3)$$

$$d + e = 4a$$

As using (4) and (5) we get that

$$a + b + c = 4a$$

Then we conclude that

$$a + b + c = d + e \tag{8}$$


Also, we have that

$$d = c(h/y) = 20a/9$$

$$e = b(h/y) = 16a/9$$

$$b = a(h/y) = 12a/9$$

And thus, we get a new triangle of the same family. This way, applying Thales theorem, we could continue constructing triangles as depicted below.



Borrallo M. On a special family of right triangles. J Pure Appl Math 2023;7(6):1.

Department of Mathematics, Nebrija University, Madrid, Spain

Correspondence: Moreno Borrallo, Department of Mathematics, Nebrija University, Madrid, Spain, E-mail: juan.morenoborrallo@gmail.com

Received: 01-Jun-2023, Manuscript No. PULJPAM-23-6489; **Editor assigned:** 05-Jun-2023, PreQC No. PULJPAM-23-6489 (PQ); **Reviewed:** 12-June-2023, QC No. PULJPAM-23-6489; **Revised:** 01-Aug-2023, Manuscript No. PULJPAM-23-6489 (R); **Published:** 08-Aug-2023, DOI: 10.37532/2752-8081.23.7(6).1.



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact reprints@pulsus.com