

# Current Research: Integrative Medicine

---

(ISSN:2529-797X)

## **Osteocalcin levels in children with metabolic syndrome: correlations with cardio-metabolic parameters**

*Maria Zhelyazkova-Savova<sup>1</sup>, Silvia Gancheva<sup>1</sup>, Sonya Galcheva<sup>2</sup>, Bistra Galunska<sup>3</sup>, Violeta Yotova<sup>2</sup>*

*<sup>1</sup>Department of Pharmacology and Clinical Pharmacology and Therapy, <sup>2</sup>Department of Pediatrics,*

*<sup>3</sup>Department of Biochemistry, Molecular Medicine and Nutrigenomics, Medical University of Varna, Bulgari*

### **Abstract**

**Background.** Osteocalcin (OC) – a bone derived vitamin K dependent protein, has been found to regulate energy metabolism. In rodents, it is the undercarboxylated form (ucOC) that has been described to release insulin from beta cells and to stimulate adiponectin secretion from adipocytes. In humans, evidence is conflicting as to which form of OC – the ucOC, or the carboxylated one (cOC) is hormonally active.

**Objective.** It was the aim of our work to study the serum levels of OC in its two forms – cOC and ucOC, and their relationships to cardio-metabolic parameters in children with metabolic syndrome.

**Methods and patients.** The population studied included 87 children - 49 boys and 38 girls (mean age  $12.06 \pm 0.32$  years, range 8-17 years). They were divided in three groups according to Cole criteria: normal (n=28), overweight (n=17) and obese (n=42). Serum levels of ucOC and cOC were measured, as were serum leptin and adiponectin levels, in addition to the conventional cardio-metabolic parameters characterizing the metabolic syndrome. Statistics: one-way ANOVA with Bonferroni multiple comparison test and the post test for linear trend were used.

**Results.** There was a clear reduction in the serum levels of cOC from normal to obese children (one-way ANOVA,  $p=0.0011$ ) with statistically significant difference between these groups, and a highly significant linear trend ( $p=0.0004$ ), while the levels of ucOC only slightly and insignificantly increased. Significant negative correlations were found between levels of cOC and BMI, body weight, waist circumference, blood glucose, HOMA-IR, triglycerides, leptin, systolic and diastolic blood pressure, and a positive correlation with adiponectin. No meaningful relationships were found between the biochemical and

anthropometric measures and ucOC.

**Conclusions.** The present results support the involvement of carboxylated OC in the regulation of carbohydrate and lipid metabolism and in the cardiovascular manifestations of metabolic syndrome at this early age.