

# Psychological Theories and Development of a Child

Maheshwari D \*

**Maheshwari D. Psychological Theories and Development of a Child. J Pedia Health Care Med 2020;3(2):1-2.**

### Abstract

Child psychology which is a broad area, covering how children change as they grow up from birth through to adolescence and trying to explain how these important changes occur. Child

psychologists attempt to make sense of every aspect of child development. Sigmund Freud gave a psychoanalytical theory explaining the psychological development of a child.

**Key Words:** *Child psychology, development, adolescence, genital stage.*

Child psychology which is a broad area, covering how children change as they grow up from birth through to adolescence and trying to explain how these important changes occur.

Child psychologists attempt to make sense of every aspect of child development, how children learn, think, interact and respond emotionally to those around them, make friends, understand emotions and their own developing personalities, temperaments and skills. [1]

### Psychological development of a Child

Sigmund Freud gave a psychoanalytical theory explaining the psychological development of a child,

He describes 6 psychosexual stages and at each stage he included the development of sexual desires. [2]

### STAGES:

#### A. Oral stage

- It is dependent stage where the infants depends on adults to fulfil his needs
- But if the child's needs are not fulfilled he develops a negative and aggressive behavior

#### B. Anal Stages

- During this stage, maturation of neuromuscular control occurs
- Child obtains a sense of control over its voluntary functions, a sense of independence.
- If it fails, child develops abnormal behavior like stubbornness.

#### C. Urethral Stage

- It is a transitional stage between anal and a phallic stage
- Child obtains and is satisfied with the control over its urinary sphincter

- If it fails to develops, child becomes competitive [3]

#### D. Phallic Stage

- Period-3rd-5th year of life
- Development of oedipus complex and electra complex where the young boys are attracted towards mother, while girls are attracted towards her father respectively.
- Child has unusual sexual feelings for the opposite sex.

#### E. Latency Stage

- This stage ends in puberty
- Child develops a well balanced control over his desires
- Child adapts to the changing environment
- Lack of this behavior results in immature behavior of the child.

#### F. Genital Stage

- Period 11-13 years to adulthood
- Child realizes sense of maturity, sense of individuality, sense of ability to reproduce, sense of independence.
- Child develops social environment by communicating and interacting with the surroundings .
- Child accepts the social expectations and lives according to it.

### CONCLUSIONS

It is therefore concluded that the behavior management is the means by which the dental health team effectively and efficiently performs dental treatment.

This can be used in both cooperative and uncooperative child.

It is the procedure which slowly develops behavior

Most often sites stents were deployed are LPA (27) and RVOT (24), raising to

challenge when stent removal in LPA and inferior vena cava (IVC) are 50% of overall locations.

---

*Malla Reddy Institute of Dental Sciences, Hyderabad, India*

*Correspondence: Maheshwari D, Malla Reddy Institute of Dental Sciences, e-mail mahi.d456@gmail.com*

*Received: July 12, 2020, Accepted: July 24, 2020, Published: July 31, 2020*



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact [reprints@pulsus.com](mailto:reprints@pulsus.com)

## Jaurena

One third of stents (28) are partially removed.

One third of stents (33) are difficult to retrieve (deep hypothermia plus circulatory arrest are requested).

50% of overall stents in LPA & RPA are partially removed.

50% of LPA & RPA stents are handled straightforward.

100% IVC stents are totally removed in deep hypothermia and decannulation.

---

## REFERENCES

1. Moffitt TE. Adolescence-limited and lifecourse-persistent antisocial behavior: a developmental taxonomy. *Psychological Review*. 1993; 100:674–701.
  2. Loeber R, Farrington DP. Young children who commit crime: epidemiology, developmental origins, risk factors, early interventions, and policy implications. *Development and Psychopathology*. 2000; 12:737–762.
  3. Snyder H. Epidemiology of official offending; in *Child Delinquents: Development, Intervention, and Service Needs*. 2001.
- 
-