

## Qualitative and quantitative screening of biochemical compositions for six selected marine macroalgae from Mediterranean coast of Egypt

Mary Guendy Naguib Ghobrial

Alexandria University, Egypt

Chemical composition of the six selected macroalgal species (*Colpomenia sinuosa*, *Padina pavonia*, *Sargassum linifolium*, *Pterocladia capillacea*, *Caulerpa racemosa*, and *Laurencia pinnatifida*) obtained from Alexandria coast of Egypt were investigated for proteins, carbohydrates, lipids, vitamins, chlorophylls, total carotenoids, and total phenols. In addition, lipidsoluble, and water-soluble antioxidant, and anti- $\alpha$ -glucosidase activities were measured for these six macroalgal species. The ash contents varied from 11.2 to 35.4 % on a dry weight basis for *P. capillacea* and *Laurencia pinnatifida*, respectively. The protein contents ranged from 5.63 % in brown macroalgae *C. sinuosa* to 8.73 % in *P. pavonia*. A relatively wide range in carbohydrate contents was observed (20.06 – 46.75 %) for the test algal species. The highest lipid percentage was found in green alga *C. racemosa* (5.91%) followed by brown algae *P. pavonia* (3.57%)

and *C. sinuosa* (2.64%). The phenolic contents varied from 1.32 mg GAE/g for *C. sinuosa* to 4.00 mg GAE/g in *P. pavonia*. The lipid-soluble compounds exhibited higher antioxidant capacity (73.18 - 145.95  $\mu$ M/g) than that of the water-soluble ones ranging from 24.83  $\mu$ M/g in *C. racemosa* to 74.07  $\mu$ M/g in *S. linifolium*. The most potent anti- $\alpha$ -glucosidase activity was observed for *P. pavonia* with IC<sub>50</sub> of 17.12  $\mu$ g/ml.

### Biography:

Mary Guendy Naguib Ghobrial was a Professor of Aquatic Plants in Hydrobiology Lab. Marine Environment Division. National Institute Of Oceanography & Fisheries (NIOF) – Ministry Of Scientific Research, Arab Republic Of Egypt (ARE). She did PhD in Botany, Faculty of Science, Alexandria University. She has published 21 research papers.