



Relapse prevention of treatment of obsessive-compulsive disorder

MS Jyotsna Singh

Department of Psychologist, India

Abstract:

Over the past three decades, obsessive-compulsive disorder (OCD) has moved from an almost untreatable, life-long psychiatric disorder to a highly manageable one. This is a very welcome change to the 1%-3% of children and adults with this disorder as, thanks to advances in both pharmacological and psychological therapies, prognosis for those afflicted with OCD is quite good in the long term, even though most have comorbid disorders that are also problematic. We still have far to go, however, until OCD can be described as either easily treatable or the effective treatments are widely known about among clinicians. This review focuses on the current state of the art in treatment for OCD and where we still are coming up short in our work as a scientific community. For example, while the impact of medications is quite strong for adults in reducing OCD symptoms, current drugs are only somewhat effective for children. In addition, there are unacceptably high relapse rates across both populations when treated with pharmacological alone. Even in the cognitive-behavioural

treatments, which show higher effect sizes and lower relapse rates than drug therapies, drop-out rates are at a quarter of those who begin treatment. This means a sizable portion of the OCD population who do obtain effective treatments (which appear to be only a portion of the overall population) are not effectively treated. Suggestions for future avenues of research are also presented. These are primarily focused on increased dissemination of effective therapies; augmentation of treatments for those with residual symptoms, both for psychotherapy and pharmacotherapy; and the impact of co morbid disorders on treatment outcome. Thirty years ago, being diagnosed with obsessive-compulsive disorder (OCD) was about the closest thing the psychiatric world had to being given a life sentence. In addition to being seen as extremely rare, prognosis for those with a diagnosis of



OCD was very poor, with no effective truly pharmacological or psychological treatments available.. Today, however, a diagnosis of OCD does not carry this loss of hope for the future and poor treatment outcomes. Instead, clinicians now have at their disposal both pharmacological and psychological treatments that are remarkably effective for the majority of patients. Still, though, there are further advances that need to be made, to continue improving treatment effectiveness and Patient outcomes. Up to 75% of persons with OCD also present with co morbid disorders.

Symptom Severity During a semi structured 1-hr interview, an independent assessor, who was a doctoral-level psychologist trained in the use of this measure, rated each patient using three 9-point Likert scales ranging from none to extreme for severity of obsessive fear, avoidance, and ritualistic behaviour.

Publication of speakers:

- 1 Franklin ME, Foa EB. Obsessive-compulsive disorder. In: Barlow DH, editor. Clinical handbook of psychological disorders. 4th ed. New York, NY: Guilford Press, 2007: 164-215
2. Lack CW, Starch EA, Murphy TK. More than just monsters under the bed: Assessing and treating pediatric OCD. *Psychiatric Times* 2006; 23: 54-57
3. American Psychiatric Association. Diagnostic and statistical 88 December 22, 2012 | www.wjgnet.com Lack CW. Research and treatments in OCD

World Psychiatrists & Psychologists Conference | August 28, 2020 | Toronto, Canada

Citation: MS Jyotsna Singh; Relapse prevention of treatment of obsessive-compulsive disorder; *Psychiatry* 2020; August 28, 2020; Toronto, Canada