Sacular aneurysm likes tibial popliteal are common, brachial artery aneurysm is rarely seen or reported in Tamilnadu, India

Kalanitthi AN, Maokar P, Thiruvengadam V

Kalanitthi AN, Maokar P, Thiruvengadam V. Sacular aneurysm likes tibial popliteal are common, brachial artery aneurysm is rarely seen or reported in Tamilnadu, India. J Vas Dis Treat. 2018;2(2):1-2.

ABSTRACT

A 60-year-old male presented with a huge mass on right arm just above the elbow joint. He had presented with these complaints with gradual progress in size over a period of 1 year. He denied any significant pain or warmth in the swelling. He also had no difficulty in elbow movements. He was seen by a general practitioner and abscess was suspected and incision and drainage was planned. Patient was examined by the author (Dr. ANK) as a second opinion. He noticed the swelling to be soft and pulsatile with no warmth or tenderness. Suspecting a vascular origin of the swelling, a vascular surgeon (Dr. VS) Consult and a CT angiogram was advised. CT Angiogram revealed a 6.1 x 5.4 x 8.1 cm sacular aneurysm of the right brachial artery with eccentric thrombus. The CT angiogram also picked up a persistent ductus arteriosus

INTRODUCTION

The patient came with a swelling in the arm near elbow joint. Patient was seen by a physician locally thought to be an abscess. Hence, he decided to incise & drain the fluid. The 60 years old gentleman reported to KHM Hospital, Anna Nagar, Chennai for second opinion. He was seen by Dr. Kalanitthi former Prof of medicine practicing in KHM Hospital. History revealed a swelling in right arm of 1 year duration gradual onset and progressively increasing in size over 15 days. On examination by Dr. AK, swelling was soft, fluctuant and pulsatile nature of the mass suspect an vascular origin. A secular aneurysm involves only a portion of the circumference resulting in an out pouching of the vessel wall (1-5).

AIMS & OBJECTIVE

To study the clinical features available and methods of treatment to be given.

MATERIALS

A single case of 60 years old gentlemen presented with a huge mass of right arm above the elbow joint. Since it was pulsatile, CT imaging done which revealed a large aneurysm, arising from the middle of the brachial artery in antero medial aspect of right upper arm with the layer of fluid and minimal edema are visualized around the aneurysm (minimal leak from the aneurysm). Incidental co existing patient ductus arteriosis was also noted during the CT imaging (6-9). The dimention was not reported in the scan center and cd was not provided hence the dimention was not given.

DISCUSSION

The opinion of vascular surgeon Dr. Vidhyasagar of Government General Hospital, Chennai was sought. He has also agreed with the physician of diagnosis of brachial artery aneurysm. After his advice patient underwent successful excision of the aneurysm and grafting. The post-operative period was uneventful and was discharged on the 3 day. Luckily patient came in

with features of pulmonary hypertension. A cardiology (Dr. PM) consult and a 2 D echo was advised. The PDA measuring 4.6 mm with left to right shunt with moderate pulmonary hypertension (PASP=46 mmHg) with mild LV Dysfunction.

Hence, a surgical excision was planned. In view of the asymptomatic cardiac status, the cardiologist (Dr. PM) advised to proceed with aneurysm excision with a moderate risk of CV events in perioperative period. He underwent successful aneurysm excision with bypass grafting and made an uneventful recovery. The case highlights the unusual location of the aneurysm hitherto unreported to literature. It also highlights the late presentation at 60 years of an asymptomatic yet hemodynamically significant PDA with such significant pulmonary hypertension.

The cardinal teaching of being beware of a pulsatile swelling is again reemphasized.

Key Words: Acute ischaemia; Embolism; Thrombosis; Arterial occlusion; Limb loss

time without undergoing incision by a local practitioner who thought to be the mass is an "abscess". Later he came to me and was shown to the reputed vascular surgeon, Dr. Vidhyasagar. It seems this case is the first case he has seen in his 30 years of practice and in my 53 years of practice curable provided by this well reputed vascular surgeon.

CONCLUSION

The case is presented due to the unique location of the aneurysm and its incidental association with a hypertensive PDA and successful treatment.

REFERENCES

- 1. Kirkov V, Iliev A, Palova DH. Variations in branching pattern of the brachial artery. Praemedicus. 2016;33(1):41-4.
- Stanchev S, Iliev A, Georgiev GP, et al. A case of bilateral variations in the arterial branching in the upper limb and clinical implications. Chronicle J Clinical Case Reports. 2017;1(1):6.
- 3. Georgiev GP, Iliev A, Dimitrova IN, et al. Palmaris longus muscle variations: significance for hand surgery and proposal of new classifications. Folia Medica. 2017;59(3):289-97.
- 4. Dimitrova IN, Trendafilova D, Iliev A, et al. Transradial catheterization failure due to high-bifurcating hypoplastic radial artery: case report. Acta Morphologica Anthropologica. 2017;24:86-8.
- 5. Ellegala DB, Day AL. Rupture of Cerebral aneurysm. N Engl J Med. 2005;352:121-4.
- 6. Patel MI, Handman DT, Fisher CM, et al. Current Views on the pathogenesis of abdominal aortic aneurysm. J Am Coll Surg. 1995;181(4):371-82.

KHM Hospital, Anna Nagar Chennai- 40, Chennai, Tamilnadu, India.

Correspondence: Dr. Kalanitthi AN, Consultant Vascular Surgeon, KHM Hospital, Anna Nagar, Chennai- 40, Chennai, Tamilnadu, India, Telephone: 9884234976, email: khmhospital1988@gmail.com

Received: March 14, 2018, Accepted: May 16, 2018, Published: May 22, 2018

This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (http:// creativecommons.org/licenses/by-nc/4.0/), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact reprints@pulsus.com

Kalanitthi et al

- 7. Bower TC, Cherry KJ, Pairolero PC. Unusual manifestations of abdominal aortic aneurysm. Surg Clin North Am. 1989;69(4):745-54.
- 8. Yeunga BKF, Pearceb WH. Surgical management of abdominal aortic aneurysm. Vas Med. 2000;5:187-93.
- 9. Burns P, Lima E, Bradbury AW. What constitutes best medicaltherapy for peripheral artery disease? Eur J Vasc Endovasc Surg. 2002;24(1):6-12.