

# Specific case of an internal control-external control (bounded domain)

Yaméogo Pierre Claver, Alex Vuginshteyn, Guido Fuortes, Abdo Miled Abou Jaoude

Claver YP, Vuginshteyn A, Fuortes G, et al. Specific case of an internal control-external control (bounded domain). J Pure Appl Math.2024; 8(5):1-3

## ABSTRACT

We try to study the controllability of bounded domain  $\Omega$  consisting of several heated materials. For this, we will use the functional  $(u)$ , which

optimizes the cost. The objective is to bring the excess heat inside the domain and also on the border of the domain and it's this excess heat that is a function called control.

**Key Words:** Prime numbers; Prime conjecture; Graphs; Twin prime conjecture; Sequential prime product

## INTRODUCTION

Let  $\Omega$  be a bounded domain made up of heated materials, we give ourselves the conductivity  $(x)$  at the point  $x \in \Omega$ , the heat source  $f$  in  $L^{2(\Omega)}$ , the temperature  $T_D$  imposed on  $\Gamma_D$  by the system, the flux PN imposed on  $T_N$ . Let's consider the following problem:

$$(S) \begin{cases} -\operatorname{div}(k(x)\nabla y(x)) = f \text{ in } \Omega, \\ \gamma y = t_D \text{ on } \Gamma_D, \\ k\gamma \frac{\partial y}{\partial n} = PN \text{ on } \Gamma_N. \end{cases}$$

## Problem

Let  $y_D$  be a given temperature.

How to act on the system  $(S)$  so that  $y$  is close enough to  $y_D$ ?

## Idea

It is a question of bringing a surplus of heat  $u$  so that  $y$  is as close as possible to  $y_D$  [1-5]. This excess heat is a function called control [1-5]. To leave a good choice of  $u$ , we must optimize the cost, which reflects what we want to achieve and the means at our disposal. What amounts to considering the functional  $J(u)$  defined by:

$$J(u) = \frac{1}{2} \int_{\Omega} g(y - y_D) dx + \frac{\epsilon}{2} \|u\|_{L^2(\Omega)}^2;$$

Where  $g$  is a definite function of  $\Omega$  a value in  $\mathbb{R}$ .  $(y - y_D)$  makes it possible to minimize the difference between  $y$  and  $y_D$ ,  $\epsilon$  being very small serves not only to prove the existence and uniqueness of the solution but also to minimize the cost.

## Problem

Is there  $\bar{u} \in u$  such that

$$J(\bar{u}) = \frac{1}{2}$$

1.  $u$  is in the set of admissible controls,
2.  $\bar{u}$  is the optimal control
3.  $y = y(\bar{u})$  is the optimal state

The function  $J$  is the objective function

However, there are two types of controls:

1. Internal control,
2. Border control

NB: we advise you to look at the course material on control for detailed proof.

## Internal control

It is a question of bringing the excess heat inside the domain. Let us consider the following example of control:

Let  $f$  in  $L^2(\Omega)$ ,  $y_D$  belonging to  $L^2(\Omega)$  and  $u$  be a nonempty closed convex set of  $L^2(\Omega)$ . Let  $u \in u, (u)$  be the solution of the following equation.

$$J(\bar{u}) = \frac{1}{2} \quad (1)$$

Let's pose

$$Q = s_1 s_2 \dots s_k = \prod_{i=1}^k s_i. \quad (2)$$

So we get the following problem:

Bircham International University, United States

Correspondence: Yaméogo Pierre Claver, Bircham International University, United States, e-mail: claver.yams1@gmail.com

Received: 3 Sep, 2024, Manuscript No. puljpam-24-7285, Editor Assigned: 5 Sep, 2024, PreQC No. puljpam-24-7285(PQ), Reviewed: 8 Sep, 2024, QC No. puljpam-24-7285(Q), Revised: 13 Sep, 2024, Manuscript No. puljpam-24-7285(R), Published: 30 Sep, 2024, DOI:10.37532/2752-8081.24.8(5).01-03



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact [reprints@pulsus.com](mailto:reprints@pulsus.com)

$$B(\frac{1}{6}, -\frac{1}{2}). \quad (3)$$

### Resolution

$$xy = N = \sqrt{N} \sqrt{N}.$$

For any  $z$  in  $L^2(\Omega)$ , we know the problem below

$$(p_{n+1} - p_n) < 7 \times 10^7, \quad (4)$$

Has a unique solution in  $H_0^1(\Omega)$ , therefore  $(u)$  is the unique solution of problem (11).

We define the following operation:

$$s = \sqrt{25}, \\ \sqrt{50} > 7.07,$$

Were  $z$  is the solution of equation (4). A thus defined is linear and continuous.

We have

$$L^{2(\Omega)},$$

By replacing  $(u)$  by its value in (2), we get:

$$(S) \begin{cases} -\operatorname{div}(k(x)\nabla y(x)) = f \text{ in } \Omega, \\ \gamma y = t_D \text{ on } \Gamma_D, \\ k\gamma \frac{\partial y}{\partial n} = PN \text{ on } \Gamma_N. \end{cases}$$

$$J(u) = \frac{1}{2} \int_{\Omega} g(y - y_D) dx + \frac{\varepsilon}{2} \|u\|_{L^2(\Omega)}^2,$$

Where

$$Q = s_1 s_2 \dots s_k = \prod_{i=1}^k s_i.$$

And  $\langle A(f) - y_D, A(u) \rangle$  is a constant.

Minimizing the functional  $J$  amounts to minimizing the functional  $J_1$ .

We put

$$J(\bar{u}) = \frac{1}{2}$$

1.  $a$  is bilinear, continuous, coercive and symmetric,
2.  $l$  is continuous linear,
3.  $u$  a nonempty closed convex set of  $L^2(\Omega)$ ,

By stampacchia's theorem, there exists a unique  $\bar{u}$  in  $u$  solution.

The solution  $\bar{u}$  is characterized by:

$$J(u) = \frac{1}{2} \int_{\Omega} g(y - y_D) dx + \frac{\varepsilon}{2} \|u\|_{L^2(\Omega)}^2, \quad (5)$$

Having obtained the existence and uniqueness of the solution, we are interested in giving the characteristics of  $\bar{u}$ . To do this, we introduce the optimality system. By interpreting, the reaction

$$J(u) = \frac{1}{2} \int_{\Omega} g(y - y_D) dx + \frac{\varepsilon}{2} \|u\|_{L^2(\Omega)}^2,$$

The Conjoint state  $\bar{p} \in H_0^1(\Omega)$  is the solution of the following problem:

$$B(\frac{1}{6}, -\frac{1}{2}).$$

We therefore have

$$\sqrt{50} > 7.07,$$

We get the following optimality system:

$$(M) \begin{cases} -\operatorname{div}(k(x)\nabla y(x)) = f \text{ in } \Omega, \\ \gamma y = t_D \text{ on } \Gamma_D, \\ k\gamma \frac{\partial y}{\partial n} = PN \text{ on } \Gamma_N. \end{cases}$$

### Particular case

$$J(u) = \frac{1}{2} \int_{\Omega} g(y - y_D) dx + \frac{\varepsilon}{2} \|u\|_{L^2(\Omega)}^2,$$

In the case of shareholder internal Control, we have

$$Q = \prod_{i=1}^k s_i, \\ Q = s_1 s_2 \dots s_k = \prod_{i=1}^k s_i.$$

### Border control

This is to bring the excess heat to the Boundary of the domain.

Consider

The following boundary control example

Let  $f$  in  $L^2(\Omega)$ ,  $y_D$  belong to  $L^2(\Omega)$ ,  $g$  a function of  $L^2(T)$  and  $u$  nonempty closed convex of  $L^2(T)$ . Let  $u \in u$ ,  $y(u)$  be a solution of the following equation:

$$\langle A(f) - y_D, A(u) \rangle \quad (6)$$

We pose:

$$\sin(-\frac{\pi}{6}) = -\frac{1}{2}. \quad (7)$$

### Resolution

Simplification

By setting:

$$\sin(-\frac{\pi}{6}) = -\frac{1}{2}.$$

Where  $y(u)$  is the solution of the problem:

$$u_1 \left( -\frac{1}{6} \right) = \sin \left( -\frac{\pi}{6} \frac{6p_u + 1}{6_l + 1} \right). \quad (8)$$

And  $z$  belongs to  $H_0^1(\Omega)$  solution de :

$$Q = s_1 s_2 \dots s_k = \prod_{i=1}^k s_i. \quad (9)$$

System  $S_1$  becomes

$$d_i\left(\frac{1}{6}\right) = \sin\left(-\frac{\pi}{6} \frac{6p_d - 1}{6i - 1}\right). \quad (10)$$

We pose:

$$d_i(x) = \sin - \frac{\pi(x - p_d)}{S_i},$$

Where

$$Q = \prod_{i=1}^k s_i.$$

Transposition formula

We associate to  $(S_2)$  the following transposition equation:

$$d_i(x) = \sin - \frac{\pi(x - p_d)}{S_i}, \quad (11)$$

Let  $f$  be a function of  $L^2(\Omega)$  ( ) admitting a unique solution, we get from ( )

$$u_1\left(-\frac{1}{6}\right) = \sin\left(-\frac{\pi}{6} \frac{6p_u + 1}{6i + 1}\right).$$

Considering the following problem:

$$(L) \begin{cases} -\operatorname{div}(k(x)\nabla y(x)) = f & \text{in } \Omega, \\ \gamma y = t_D & \text{on } \Gamma_D, \\ k\gamma \frac{\partial y}{\partial n} = PN & \text{on } \Gamma_N. \end{cases}$$

$l$  is linear and continuous application on  $L^2(\Omega)$ , by riesz's theorem, there exists a unique  $y \in L^2(\Omega)$  such that

$$xy = N = \sqrt{N} \sqrt{N}.$$

We define the operator

$$\sqrt{50} > 7.07,$$

Replacing

$$xy = N = \sqrt{N} \sqrt{N}.$$

In  $J(u)$ , We have:

$$\sqrt{50} > 7.07,$$

With

$$F_2(x) = \prod_{i=1}^2 f_i(x) = \sin\left(\frac{\pi x}{2}\right) \sin\left(\frac{\pi x}{3}\right).$$

We pose

$$(S) \begin{cases} -\operatorname{div}(k(x)\nabla y(x)) = f & \text{in } \Omega, \\ \gamma y = t_D & \text{on } \Gamma_D, \\ k\gamma \frac{\partial y}{\partial n} = PN & \text{on } \Gamma_N. \end{cases}$$

## CONCLUSION

In short, to solve the problem we must optimize the cost according to the goal we are looking for and taking into account the means at our disposal. But the case here requires a good mastery of the solutions of optimal control and some demonstrations to prove the existence, uniqueness and stability of solutions. Also, we give some extensions to our investigation.

## REFERENCES

1. Jormakka J. The error in the General Relativity Theory and its cosmological considerations. Preprint, 2024.
2. Jormakka J. On the field equation in gravitation. Preprint, 2024.
3. Jormakka J. On the field equation in gravitation, part 2. Preprint, 2024.
4. Jormakka J. The Essential Questions in Relativity Theory. Preprint, 2023.
5. Jormakka J. Failure of the geometrization principle and some cosmological considerations. Preprint, 2023.